

This proposal should be viewed in color. If your hardcopy is not in color, please request a color copy.


## SMOKE SIGNALS FROM OPHIUCHUS

Alyssa Goodman, Bryan Gaensler, Charles Lada, Scott Schnee & Scott Wolk

*Harvard-Smithsonian Center for Astrophysics*

This proposal offers XMM-Newton a chance to solve a mystery. The composite figure below, explained in this proposal, takes us through the first many chapters of an obviously exciting story—all about giant rings of smoke being blown into the interstellar medium. When completed, the story will tell a great tale of many generations of stars forming in Ophiuchus. Trouble is, without some help from XMM-Newton, we still don't know much about the protagonist(s).

**Figure 1**



**In each panel where it is shown, the white ring shows a 2 pc circle, corresponding to the size and shape of the heated ring apparent in the IRAS Temperature Map.**


This proposal should be viewed in color. If your hardcopy is not in color, please request a color copy.

### Background Information: Will the Real $\rho$ -Oph Cluster Please Stand Up?

Ophiuchus is one of a handful of well-studied star-forming regions within 200 pc of the Sun. The distance to Ophiuchus is  $\sim 160$  pc, so the 1-degree grid boxes shown in Figure 1 correspond to about 2 pc in our region of interest. The area labeled “Region known as  $\rho$ -Oph Cluster” in Figure 1 contains more than 400 forming stars, the most massive of which is a B star (Bontemps et al. 2001; Wilking, Lada & Young 1989). Trouble is, it turns out this region is only known as the “ $\rho$ -Oph Cluster” because it is *near* the young BIV star  $\rho$ -Oph—not around it (see the labeled yellow star in Figure 1’s lower right panel). Now, though, thanks to a ring of warm smoke revealed by new IRAS maps and a gaggle of embedded sources revealed by ROSAT (see Figure 1), we have found the *real*  $\rho$ -Oph Cluster, and we would like to study it<sup>1</sup>.

The new IRAS-based column density and temperature maps shown in Figure 1 (from Goodman, Li & Schnee 2003) were created in conjunction with the ongoing **CO**ordinated **M**olecular **P**robe **L**ine **E**xinction **T**hermal **E**mission (**COMPLETE**) Survey of Star Forming Regions. COMPLETE is a large, international, coordinated effort to map three  $>10$ -pc-scale star forming regions being mapped under the SIRTf Legacy Cores-to-Disks (c2d) program with every relevant ground-based method of measuring density, temperature and velocity structure.<sup>2</sup> In order to create the new IRAS maps, we re-calibrated the zero points of the IRAS Sky Survey Atlas (ISSA) 60 and 100 micron plates, to facilitate their use down to very low flux levels.

Upon examining the “new & improved” IRAS maps made for COMPLETE, we *discovered* the 2-pc diameter heated dust ring shown in Figures 1 and 2. The temperature of that ring is only about 37 K, or 7 K hotter than its immediate environment. After further investigation, we determined that the unusual X-ray source 1RXSJ162554.5-233037 (discussed below) lies very near the ring’s center. We also noticed that the elevated-temperature ring fits neatly inside a crook of dust emission extending North from the  $\rho$ -Oph region (see Figure 1).



**Figure 2:** *Left panel:* Nebulosity around  $\rho$ -Oph (DSS); *Right Panel:* IRAS temperature image of shell (colors from 26 to 39 K only). The **maroon ring** shows the same 2 pc diameter shell as on the previous page, and the **green star** shows 1RXS J162554-233037.  $\rho$ -Oph is apparent in the left panel with diffraction spikes.

With great surprise, we also realized that the ring we detected in IRAS encircles the famous star  $\rho$ -Ophiuchi, for which the (famous, but fraudulently named) “ $\rho$ -Oph cluster” a full degree south of (the real)  $\rho$ -Oph is named.  $\rho$ -Oph itself is a BIV star, and it powers an oft-photographed very beautiful bright nebula, shown in black and white in Figure 2. The existence of this nebula historically made it difficult to search in the optical for a cluster *really* associated with  $\rho$ -Oph, and *until now, none was known.*

Fortuitously, the region around (the real)  $\rho$ -Oph, interior to the IRAS ring, was serendipitously included in a ROSAT pointed observation made by Snowden et al. in 1994. The data from the pointed observation show about twenty ROSAT-detected point sources within the IRAS ring, all with heavily “reddened” hardness ratios,

suggestive that the sources form a loose embedded cluster associated with  $\rho$ -Oph. One source, **1RXSJ162554.5-233037** has a well-determined Hardness Ratio 2=1.0, and Hardness Ratio 1=0.6, making it an excellent candidate to be some kind of “exotic” compact object (e.g. neutron star, accreting binary). Furthermore, with only a small dose of imagination, one can see that most of the ROSAT sources discovered around  $\rho$ -Oph lie on another ring, interior to the IRAS ring (see the light dashed line in Figure 1). The XMM-Newton observations proposed below, in concert with complementary Chandra observations proposed last month (and described below), will easily reveal whether or not this ring in the pattern the ROSAT sources was an illusion or not.

<sup>1</sup> Do not be fooled by papers in print with titles like “X-rays and regions of star formation: a combined ROSAT-HRI/ near-to-mid IR study of the rho Oph dark cloud” (Grosso et al. 2000) or “CHANDRA Observations of the  $\rho$ -Ophiuchi Cloud” (Imanishi et al. 2001). Every targeted X-ray observation of this region to date has been of the southern, well-known cluster, not the one around the “real”  $\rho$ -Oph.

<sup>2</sup> Please see <http://cfa-www.harvard.edu/COMPLETE> for more information.

*This proposal should be viewed in color. If your hardcopy is not in color, please request a color copy.*

### *What caused the ring in Ophiuchus, and what might it have to do with sequential star formation?*

The most obvious astronomical hypothesis when a warm nearly-perfect circle is found potentially centered on an exotic X-ray source is that a powerful wind, maybe even a supernova, created a shell in the interstellar medium which we observe as a limb-brightened ring.

To evaluate the plausibility of the Ophiuchus ring being a supernova, we can use the simple formalism set out in Mike Shull's "Signature of a Buried Supernova" paper of 1980, to calculate that a  $0.5 \times 10^{51}$  erg supernova ejecting a mass of 1 solar mass into a medium, like the dense parts of Ophiuchus, with a density of  $10^5 \text{ cm}^{-3}$  would reach the size apparent in the Figures ( $\sim 2$  pc) in about 200,000 years. At that time, the temperature of the shell would be 38 K—eerily like the color temperature we observe. The velocity of this shell at 200,000 years would be only 1.7 km  $\text{s}^{-1}$ , which is only slightly larger than the  $^{13}\text{CO}$  line widths in this region, and thus nearly undetectable as an expansion—unless one were explicitly looking for such a signature. The (well-known, southern) "[Oph]" cluster contains some stars as old as  $\sim 1$  Myr, so we do *not* think that this hypothetical 200,000-year-old supernova would have directly "triggered" the formation of the (well-known, southern) "[Oph]" cluster—but the calculation certainly makes one wonder whether the [Oph] molecular cloud complex has at least been struck by at least one supernova since it began forming stars.<sup>3</sup> Also, the (possibly accidental) ring-like pattern of the ROSAT sources apparent in Figure 1, which is concentric with and interior to the IRAS ring, leads one to dream that perhaps this hypothetical supernova could even have "exploded" its natal cluster, sending forming stars to the (constant) radial distance from the explosion's center where we see them now.

But, SNe are not the only way to drive powerful winds into the ISM—a substantial wind should be associated with the B star [Oph] itself. At this time, though, without more information on the sources around (the real) [Oph], and more information about the mass<sup>4</sup> and velocity of the ring, it is hard for us to evaluate the host of possible hypotheses. *Identifying the creator of the IRAS ring, and constraining its former identity (SN progenitor, star with wind, accreting binary, etc.) is a key goal of our proposed observations.*

Our longer-term goal, which also justifies much of the COMPLETE Survey, is to *evaluate the role of previous generations of stars on the long-term productivity of star-forming molecular gas*. For example, with COMPLETE and SIRTf data, we will be able to use molecular-line observations along with a point-source census to statistically evaluate the role of pc-scale bipolar outflows from young stars in churning and destroying their natal clouds. By studying large shells like the one described in this proposal, we are investigating the influence of prior generations of stars on their associated molecular clouds on very large ( $\gg 1$  pc) spatial scales, and honestly unknown timescales. Imagine, for example, that star-formation in one region (e.g. the southern well-known "[Oph] cluster") is accelerated by increased pressure caused by the expansion of a neighboring shell (e.g. the one associated with 1RXSJ162554.5-233037). This hypothesis, similar to the one put forward as "sequential star formation" by Elmegreen and Lada (1977), implies that one *cannot* legitimately calculate star-formation rates and/or efficiencies, in our Galaxy or anywhere else, without taking environmental effects—on large scales—into account.

### *Clues from all wavelengths*

The missing chapters of the story we described at the outset tell what protagonist(s) caused the warm dust ring around 1RXSJ162554.5-233037, and how and when the shell was created. The research for these missing chapters necessitates a multi-wavelength approach. Existing radio continuum data are insufficiently sensitive to find the synchrotron emission that might be associated with this shell, so we are in the process of proposing to search for it later this year. Radio-wavelength molecular or HI-absorption spectral-line data might reveal expansion in the shell, and we are in the process, as part of COMPLETE, of acquiring the relevant data at FCRAO and the GBT. Far-IR data gave us the introductory chapter to our mystery, by revealing the Ophiuchus ring. Existing (just released!) 2MASS and upcoming SIRTf near-IR data will give much needed information about the age and mass distribution of the cluster associated with (the real) [Oph], as well as the extinction structure within the shell.<sup>5</sup> The near-IR data are critical, since optical images of this region are dominated by the nebulosity around [Oph]. And, last but

---

<sup>3</sup> On the longer time scale, and larger spatial scale, de Geus and colleagues argued very convincingly that the bubbly and filamentary nature of the entire Sco-Cen-Oph region of the ISM can be explained as due to the effects of stellar associations and their winds upon each other (de Geus 1992; de Geus, Bronfman, & Thaddeus 1990; de Geus & Burton 1991; de Geus, de Zeeuw, & Lub 1989).

<sup>4</sup> As part of COMPLETE, we have been awarded a large amount of time on the JCMT, where we are using the SCUBA instrument to map out dust mass. The region of the IRAS ring in Ophiuchus was observed in March 2003, and the data are currently being reduced.

<sup>5</sup> As part of COMPLETE, we are using a sophisticated star-counting algorithm known as "NICER" to map extinction without the bias caused by dust temperature when using thermal dust emission measurements (see Lombardi & Alves 2001 and references therein).

*This proposal should be viewed in color. If your hardcopy is not in color, please request a color copy.*

certainly not least, X-rays may well hold the key to unlocking the mystery. The ROSAT data have already strongly hinted that 1RXSJ162554.5-233037 may well be our protagonist(s), but we need to know much more about this character before writing more of our story.

### *Observations Planned with XMM-Newton and Chandra*

While our overall goal is to understand what created the newly-discovered dust shell shown in the Figures, the goal of our proposed X-ray observations are these:

- 1) Determine the spectral nature of the unusual source at the center of the cluster.
- 2) Determine the nature of the X-ray sources within the cluster to determine if they are (young) stars.
- 3) Understand the nature of the objects in the “X-ray ring.”

Our plan for Chandra-HRI (proposed last month) is to map enough of the (real)  $\square$ -Oph cluster region to accurately determine the positions of all cluster members, while simultaneously covering a large enough area to survey nearly all of the ring discovered in the new IRAS temperature maps. HRI on Chandra gives us the wide field required, but no spectral information will come from those observations, and count rates will be low. We propose to use XMM-Newton to determine the spectral nature of the brighter sources we will map out with Chandra through a set of three overlapping 15ks observations. The XMM-Newton visibility checker shows no problems with observing this region of Ophiuchus in this way. Each proposed pointing of XMM-Newton is represented as a green circle in Figure 1’s lower-right panel, and the four proposed pointings of Chandra-HRI are shown as white-dotted squares.

If 1RXSJ162554.5-233037 is as exotic as the ROSAT observations suggest it may be, the spectra will give us insight in to what kind of source it is. ROSAT saw a count rate in 1RXSJ162554.5-233037 of about 0.016 cps. Using PIMMS we modeled this source as a  $\square=2$  power law with  $N_{\text{H}}=10^{22}$   $\text{cm}^{-2}$ . From this we estimate a count rate of 0.76 in the PN with the medium filter and 1.33 cps in the sum of the EPIC+MOS detectors. We fit the simulated spectrum of the central object and were able to recover the power-law and column to less than 5%. Iron emission will be easy to detect if present. Pileup will be small, only about 4%. Most of the sources in the ROSAT field are as bright as 1RXSJ162554.5-233037 so similar quality spectra will be obtained. The time resolution of the PN detector gives us sensitivity to periods slower than 150 ms, appropriate to detect all but the very youngest (<10 kyr) rotation-powered pulsars. With about 10,000 counts per observation in the PN detector and assuming a sinusoidal pulse profile, a blind periodicity search will be sensitive to pulsed fractions as small as 10%. We could improve this by another factor of  $\sim 2$  if the pulse shape is narrow or if the three observations are closely spaced in time.

The key argument for the depth of our proposed exposure is not the sources we do see (in the ROSAT data), but the sources we do not see. We suspect that the region interior to the warm dust ring in Ophiuchus is home to a previously unknown stellar cluster centered about  $\square$  Oph, and that a member of this cluster (possibly 1RXSJ162554.5-233037) has created the dust ring currently seen to be impacting the Ophiuchus molecular clouds (see Figure 1). If the cluster were very young, one would expect to see near-infrared excesses consistent with the presence of circumstellar disks around the members. An analysis of near-IR 2MASS data shows no such excesses, implying that this is a cluster of “older,” diskless, pre-main-sequence stars. Several studies have shown that the best tracer of diskless PMS stars is X-ray emission, and only X-ray telescopes with good sensitivity at hard energies can identify PMS stars in regions of obscuration (Feigelson & Montmerle 1999). In a 15 ks observation we will obtain about 100 counts from a source with an unabsorbed flux of about  $10^{-13}$   $\text{ergs cm}^{-2} \text{sec}^{-1}$  or  $L_x=10^{29}$  at the distance of the cloud ( $\sim 0.35M_{\odot}$ ). The expected count rates for cluster members will allow us to determine each member’s absorbing column ( $N_{\text{H}}$ ), rough temperature, and approximate age when we combine X-ray and 2MASS data (giving  $L_x/L_{\text{bol}}$ ).

As shown in Figure 1, the three pointings requested here cover most of the region we plan to map out with Chandra-HRI. The pointings are designed to optimally sample the central region of the dust ring, giving maximum signal for 1RXSJ162554.5-233037 while still covering most of the “ring” of ROSAT sources. Note that due to the high brightness of  $\square$ -Oph itself, we have not requested use of the Optical Monitor in this proposal.

### *References*

- Bontemps, S. et al. 2001, A&A, 372, 173  
de Geus, E.J. 1992, A&A, 262, 258  
de Geus, E.J., Bronfman, L., & Thaddeus, P. 1990, A&A, 231, 137  
de Geus, E.J., de Zeeuw, P.T., & Lub, J. 1989, A&A, 216, 44  
Elmegreen, B.G. & Lada, C.J. 1977, ApJ, 214, 725  
Feigelson, E. & Montmerle, T. 1999, ARAA.  
Goodman, A.A., Li, J.G. & Schnee, S. 2003, ApJL, in prep.  
Grosso, N., Montmerle, T., Bontemps, S., André, P., & Feigelson, E. D. 2000, A&A, 359, 113  
Imanishi, K., Koyama, K., & Tsuboi, Y. 2001, ApJ, 557, 747  
Lombardi, M., & Alves, J. 2001, A&A, 377, 1023  
Wilking, B.A., Lada, C.J., & Young, E.T. 1989, ApJ, 340, 823